



## THE EVALUATION OF THE CONNECTION BETWEEN MOTOR PERFORMANCE SKILLS AND BODY COMPOSITION OF THE GIRLS

Nurten Dinç,  
Mehmet Hilmi Gökmen<sup>i</sup>,  
Murat Taş,  
Burcu Kıvrak,  
Canberk Çiçekli

Manisa Celal Bayar University,  
Coaching Training Department,  
Manisa, Turkey

### Abstract:

**The Purpose:** This study was done for evaluating the connection between motor performance skills and body composition of the girls between 6-10 ages. **The Method:** 57 schoolgirls between 6-10 ages were included to the study. Weight, Age, Height, Body Mass Index (BMI), Free Fat Mass (FFM), Lean Muscle Mass (LMM), Body Fat Mass (BFM) and Body Fat Percentage (BFP) values were measured with (Inbody 230) branded tanita. Long jump, right hand and left hand strength, back strength, 20 meters speed test, sit & reach test, 1 minute pull-up and flamingo balance tests were applied to the participants. **Results:** Pearson Correlation Test was applied for the analysis by using SPSS 23.0 program. According to the results, the average age is (8,00±,926) years, the average height is (128,7±6,64) cm and the average weight is (28,6±6,67) kg. There is a statistically meaningful relation between long jump, right hand, left hand, back strength, flamingo balance test and BFM, FFM, LMM, BMI and BFP at the (p<0.01) level. However, there is no relation found between 20 meters speed test, sit & reach test, 1 minute pull-up test and body composition parameters. **Discussion and Conclusion:** At the end of our study, we can say that there is a meaningful relation in a positive way between some motor performance skills and body composition of the girls.

**Keywords:** body composition, girls, balance, flexibility

---

<sup>i</sup> Correspondence: email [mehmethilmi93@gmail.com](mailto:mehmethilmi93@gmail.com)

## 1. Introduction

For last 20-30 years, physical activity levels of the children have changed dramatically. Besides, they are spending more time in indoor places over outdoor places for their physical activities. They go to the school by bus or by car instead of by foot or by bicycle. This situation leads low rates for doing exercises (Fjørtoft et al. 2011). Health problems mostly seen in elderly ages like overweighting (obesity), cardiovascular diseases, diabetes, musculoskeletal diseases cause low standard of living and many risks. On these days, these diseases are seen in teenage periods (Haslofça et al. 2011). In this context, motor performance skills can be considered as an important matter for physical suitability and motor development levels of the children. The children who have better motor performance skills are more physically active and have lower chance of being inactive in contrast to the children who have worse motor performance skills. Also, it's a fact that in order to improve the motor performance skills of the children, activities like running, pace, agility, jumping and balance are needed to be applied (Karim et al. 2015).

Basic movement skills are observed fundamental action forms of the childhood. Movements during sports, exercise and physical activity include physical skills like running, bouncing, jumping, balance, contort, escaping, throwing and catching (Brien et al. 2016). Gaining basic movement skills (BMS) are listed developmentally (biological, psychological, sociological, motivational, cognitive etc.). The process of gaining these skills contains active play experiences and created programs (Siahkouhian et al. 2011). As BMS sufficiencies increase, health and physical sufficiency increase. Besides, ideal body weight maintaining rate grows (Bryant et al. 2014). According to some results, BMS is related to physical activity level. Not improving these skills in childhood may affect physical activity level in adulthood (Comeau et al. 2017). For example, Henrique and his friends have made a research on 293 children. After 2 years of monitoring physical activity, they found that BMS are improved in the ratio of 21% (Henrique et al. 2016). In the guidance of these knowledgements, our study was made for evaluating motor performance skills and body composition relations of the girls.

## 2. Material and Method

The study was applied in İzmir Özel Türk Koleji, Bornova. 57 schoolgirls have attended to the study with (8,00±,926) years average age, (128,7±6,64) cm average height and (28,6±6,67) kg average weight. The study period was in their school time which is between 9 am and 4 pm. Before the study, their parents have signed papers for volunteering. These tests were applied to the girls in order. Body Composition Measurement, Long Jump Test, Right Hand and Left Hand Strength Test, Back Strength, 1 Minute Pull-Up Test, 20 Meters Speed Test, Sit & Reach Test and Flamingo Balance Test.

## 2.1 Statistical Analysis

The data analyze was made with SPSS 23.0 program. Frequency analyze was made to find the height, weight and age averages of children. Pearson Correlation Test was applied in order to specify the relationship between their performance tests and body composition measurements.

## 3. Results and Discussion

**Table 1:** Average and Standard Deviation Values of the Participants

Variables	N	Average (X) ± SS
Age (years)	57	8,00±,926
Height (cm)	57	128,7±6,64
Body Weight (kg)	57	28,6±6,67
BMI (kg/m <sup>2</sup> )	57	17,1±2,73
BFP (%)	57	24,0±7,84
FFM (%)	57	21,3±3,24
BFM (kg)	57	7,32±4,03
LMM (kg)	57	10,5±1,93
Long Jump (cm)	57	122,3±14,5
Right Hand Strength (kg)	57	10,7±2,60
Left Hand Strength (kg)	57	9,73±2,30
Back Strength (kg)	57	28,7±6,20
20 meters Speed Test (sn.)	57	5,19±,623
Sit & Reach Test (cm)	57	10,6±4,63
1 min Pull-Up Test (number)	57	21,1±4,16
Flamingo Balance Test (n)	57	16,4±8,28

Average and standard deviation values of physical parameters and performance tests of the participants were given in Table 1.

**Table 2:** Relation between Body Composition and Motor Performance Skills of the Girls

Variables	Long Jump	Right Hand Strength	Left Hand Strength	Back Strength	20 meters Speed Test	Sit & Reach Test	1 min Pull-Up Test	Flamingo Balance Test
<b>Long Jump</b>	1	,386**	,361**	,303*	-,536**	,361**	,471**	,034
Right Hand Strength	,386**	1	,852**	,633**	-,307*	,203	,210	,264*
Left Hand Strength	,361**	,852**	1	,703**	-,308*	,199	,199	,275*
Back Strength	,303*	,633**	,703**	1	-,308*	,239	,324*	,128
20 meters Speed Test	-,536**	-,307*	-,308*	-,308*	1	-,306*	-,457**	,015
Sit & Reach Test	,361**	,203	,199	,239	-,306*	1	,188	-,210
1 min Pull-Up Test	,471**	,210	,199	,324*	-,457**	,188	1	,058

Nurten Dinç, Mehmet Hilmi Gökmen, Murat Taş, Burcu Kıvrak, Canberk Çiçekli  
THE EVALUATION OF THE CONNECTION BETWEEN MOTOR PERFORMANCE SKILLS AND  
BODY COMPOSITION OF THE GIRLS

Flamingo Balance Test	,034	<b>,264*</b>	<b>,275*</b>	,128	,015	-,210	,058	1
BFM	,103	<b>,445**</b>	<b>,491**</b>	<b>,326*</b>	,006	-,040	-,060	,229
FFM	<b>,312*</b>	<b>,690**</b>	<b>,750**</b>	<b>,557**</b>	-,160	-,101	,073	<b>,369**</b>
LMM	<b>,325*</b>	<b>,685**</b>	<b>,749**</b>	<b>,561**</b>	-,161	-,095	,081	<b>,373**</b>
BMI	,095	<b>,444**</b>	<b>,510**</b>	,325	,035	-,003	-,074	,260
BFP	,003	<b>,275**</b>	<b>,302*</b>	,164	,092	-,101	-,080	,171

\*P< 0.05; \*\*P< 0.01

There is a meaningful difference between Long Jump Test and Right Hand Strength ( $r=,386$ ;  $p<0.01$ ), Left Hand Strength ( $r=,361$ ;  $p<0.01$ ), Back Strength ( $r=,303$ ;  $P<0.05$ ), 20 Meters Speed Test ( $r= -,536$ ;  $p<0.01$ ), Sit & Reach Test ( $r=,361$ ;  $p<0.01$ ), 1 Min. Pull-Up Test ( $r=,471$ ;  $p<0.01$ ), Free Fat Mass (FFM) ( $r=,312$ ;  $p<0.05$ ) and Lean Muscle Mass (LMM) ( $r=,325$ ;  $p<0.05$ ) levels. However, there isn't a meaningful difference between Flamingo Balance Test, Body Fat Mass (BFM), Body Mass Index (BMI) and Body Fat Percentage (BFP).

There is a meaningful difference between Right Hand Strength and Left Hand Strength ( $r=,852$ ;  $p<0.01$ ), Back Strength ( $r=,633$ ;  $P<0.01$ ), 20 Meters Speed Test ( $r= -,307$ ;  $p<0.05$ ), Flamingo Balance Test ( $r=,264$ ;  $p<0.05$ ), BFM ( $r=,445$ ;  $p<0.01$ ), FFM ( $r=,690$ ;  $p<0.01$ ), LMM ( $r=,685$ ;  $p<0.01$ ), BMI ( $r=,444$ ;  $p<0.01$ ) and BFP ( $r=,275$ ;  $p<0.01$ ) levels, while there isn't a meaningful difference between Sit & Reach Test and 1 Min. Pull-Up Test.

A meaningful difference between girls' Left Hand Strength and Back Strength ( $r=,703$ ;  $P<0.01$ ), 20 Meters Speed Test ( $r= -,308$ ;  $p<0.05$ ), Flamingo Balance Test ( $r=,275$ ;  $p<0.05$ ), BFM ( $r=,491$ ;  $p<0.01$ ), FFM ( $r=,750$ ;  $p<0.01$ ), LMM ( $r=,749$ ;  $p<0.01$ ), BMI ( $r=,510$ ;  $p<0.01$ ) ve BFP ( $r=,302$ ;  $p<0.05$ ) levels but no meaningful difference is found between Sit & Reach Test and 1 Min. Pull-Up Test.

There is a meaningful difference between Back Strength and 20 Meters Speed Test ( $r= -,308$ ;  $p<0.05$ ), 1 Min. Pull-Up Test ( $r=,324$ ;  $p<0.05$ ), BFM ( $r=,326$ ;  $p<0.05$ ), FFM ( $r=,577$ ;  $p<0.01$ ), LMM ( $r=,561$ ;  $p<0.01$ ) and BMI ( $r=,325$ ;  $p<0.05$ ) levels but there isn't a meaningful difference between Sit & Reach Test, Flamingo Balance Test and BFP.

A meaningful difference is found between 20 Meters Speed Test and Sit & Reach Test ( $r= -,306$ ;  $p<0.05$ ), 1 Min. Pull-Up Test ( $r= -,457$ ;  $p<0.05$ ) levels. However, there isn't a meaningful difference between Flamingo Balance Test, BFM, FFM, LMM, BMI and BFP.

There isn't a meaningful difference found between Sit & Reach Test and Right Hand Strength, Left Hand Strength, Back Strength, 1 Min. Pull-Up Test, Flamingo Balance Test, BFM, FFM, LMM, BMI and BFP.

There isn't a meaningful difference between girls' 1 Min. Pull-Up Test and Right Hand Strength Test, Left Hand Strength Test, Sit & Reach Test, Flamingo Balance Test, BFM, FFM, LMM, BMI and BFP.

A meaningful difference is observed between Flamingo Balance Test and FFM ( $r=.369$ ;  $p<0.01$ ) levels while there's not a meaningful difference between Long Jump Test, Back Strength, 20 Meters Speed Test, Sit & Reach Test, 1 Min. Pull-Up Test, BFM, BMI and BFP.

According to our study, BFP average of the girls is  $24,0\pm 7,84$ . BFP average of the girls is  $23,4\pm 7,7$  as a result of the study that Marmeleira et al. (2017) have made about the relations between motor performance and body compositions of the children between 6-10 ages. Moreover, Ozkocak (2018) made a study on body composition of the children between 5-14 years and found that BFP average of the girls is  $24,77\pm 6,31$ . The results in Marmeleira and friends (2017) and Ozkocak (2018) are close to our study findings.

Our study has showed that BMI average of the girls is  $17,1\pm 2,7$ . According to the study on the children between 6-10 ages, Souza et al. (2014) have found that BMI average of the girls is  $16,16\pm 1,47$ . Duncan et al. (2017) made a study about relationship between BMI average and basic motor skills of the children between 6-11 years and they found that BMI averages of the girls is  $16,7\pm 0,3$ . Santos et al. (2014) made a study about relationship between body composition and motor performance of the children between 7-10 years and they found that BMI average of the girls is  $17,0\pm 2,9$ . In addition, Kim et al. (2016) made a search on pre-school children in Korea and they found that BMI average of the girls is  $16,3\pm 0,62$ . As we made a research on other studies in the literature, the results are close to our study findings.

We found the average 20 meters Speed Test of the girls is  $5,19\pm 0,62$ . Román et al. (2016) made a search on pre-school children and observed that the average 20 meters Speed Test of the girls is  $6,25\pm 1,16$ . Dumit et al. (2010) studied about obesity and physical suitability on the children and adolescents. They found the average 20 meters Speed Test of the girls is  $4,40\pm 0,40$ . Moreover, Sanchez et al. (2015) made an analyze on pre-school children for physical suitability and the average 20 meters Speed Test of the girls is found as  $5,80\pm 0,71$ . Also, Roman et al. (2015) made a study about identifying physical suitability of the children between 3-6 ages and found the average 20 meters Speed Test of the girls is  $6,37\pm 1,08$ . Our findings are similar as other literature findings.

When we analyzed the relation between BMI and both hands strength of the girls, there is a ( $p<0.01$ ) level statistically meaningful relation. In addition to our study, the findings are similar as Montalcini et al. (2016), Souza et al. (2014), Liao (2016) and Ervin et al. (2014) results and Lad et al. (2013) study result is ( $P<0.05$ ) level. However, Omar et al. (2014) findings are opposite to our findings.

In our study, the average Sit & Reach Test of the girls is found as  $16,4\pm 8,28$ . Mayorga et al. (2013) made a study on the children between 10-12 ages. They found that the average Sit & Reach Test of the girls is  $30,30\pm 6,34$ . On the other hand, Chillón et al. (2011) have made a study about contrasting the Spanish children and adolescents. They found the average Sit & Reach Test of the girls is  $19,4\pm 5,5$ . Celik et al. (2013) tested motoric features of 7-9 age group children and observed the average Sit & Reach Test of

the girls is  $18,72 \pm 4,65$ . Urlu (2014) analyzed physical activity level of the children between 10-12 years old and found the average Sit & Reach Test of the girls that study on 4<sup>th</sup> semester is  $20,87 \pm 7,36$ . Casonatto et al. (2015) made a study about physical suitability and BMI of the children and observed the average Sit & Reach Test of the girls is  $29,0 \pm 8,0$ . Chaves et al. (2016) analyzed Peruvian children between 6-14 ages for motor improvements. They found the average Sit & Reach Test of the 10 years old girls group is  $22,1 \pm 1,3$ . The results are similar as Mayorga et al. (2013), Chillón et al. (2011), Celik et al. (2013), Urlu (2014) and Chaves et al. (2016) studies but Casonatto et al. (2015) study findings are higher than ours so there is an opposition with our findings.

Long Jump Test average of the children is found as  $122,3 \pm 14,5$ . Addition to our studies, Santos et al. (2015) made a research about lower body strength of the children and they found Long Jump Test average of the girls is  $127,35 \pm 20,3$ . Ranson et al. (2015) studied on school children and they found Long Jump Test average of the children is  $128,97 \pm 19,29$ . Zaqout et al. (2016) made a study about physical suitability of the European children and observed that Long Jump Test average of the girls is  $110,9 \pm 24,0$ . Henrique et al. (2017) analyzed motor improvement performances of the Portuguese children and found Long Jump Test average of the 9 years old girls is  $104,7 \pm 38,5$ . Lastly, Chaves et al. (2015) analyzed motor improvements of the Portuguese children between 6-10 years and they observed Long Jump Test average of the girls is  $111,60 \pm 20,32$ . Our results are same as Zaqout et al. (2016), Henrique et al. (2017) and Chaves et al. (2015). On the other hand, results in Santos et al. (2015) and Ranson et al. (2015) examinations are higher than our findings.

#### 4. Conclusion

In conclusion, we can say that there is a meaningful connection in positive way between motor performance skills and body composition of the girls.

#### About the Author(s)

**Nurten Dinç** is Assistant Professor, Department of Sports Health, Manisa Celal Bayar University.

**Mehmet Hilmi Gökmen** is Master Student Department of Sports Health, Manisa Celal Bayar University.

**Murat Taş** is Professor, Department of Sports Health, Manisa Celal Bayar University.

**Burcu Kıvrak** is Master Student Department of Sports Health, Manisa Celal Bayar University.

**Canberk Çiçekli** is Master Student Department of Sports Health, Manisa Celal Bayar University.

## References

- Brien W.O., Belton S., Issartel J., 2016. The relationship between adolescents' physical activity, fundamental movement skills and weight status. *Journal of Sports Sciences*. 12; 1159–1167. <http://dx.doi.org/10.1080/02640414.2015.1096017>.
- Bryant E.S., Duncan M.J., Birch S.L., 2014. Fundamental movement skills and weight status in British primary school children. *European Journal of Sport Science* 7; 730–736. <https://doi.org/10.1080/17461391.2013.870232>.
- Casonatto J., Fernandes R.A., Batista M.A., Cyrino E.S., Silva M.J., 2015, Arruda M.D., Ronque E.R. Association between health-related physical fitness and body mass index status in children. *Journal of Child Health Care* 1-10. doi: 10.1177/1367493515598645.
- Chaves R., Jones A.B., Gomes T., Souza M, Pereira S, Maia J, 2015. Effects of individual and school-level characteristics on a child's gross motor coordination development. *International Journal of Environmental Research and Public Health*. 12; 8883-8896. doi:10.3390/ijerph120808883.
- Chaves R.N.D., Valdúvia A.B., Nevill A., Freitas D., Tani G., Katzmarzyk P.T., Maia J.A.R., 2016. Developmental and physical-fitness associations with gross motor coordination problems in Peruvian children. *Research in Developmental Disabilities*. 54; 107- 114. <http://dx.doi.org/10.1016/j.ridd.2016.01.003>.
- Chillón P., Ortega F.B., Ferrando J.A., Casajus J.A., 2011. Physical fitness in rural and urban children and adolescents from Spain. *Journal of Science and Medicine in Sport* 14; 417-423. doi:10.1016/j.jsams.2011.04.004.
- Comeau M.E., Bouchar, D.R., Levesque C., Johnson M.J., Rioux B.V., Mayo A., Sénéchal M., 2017. Association between functional movements skills and health indicators in children aged between 9 and 12 years old. *Int. J. Environ. Res. Public Health* 14, 1-15. doi:10.3390/ijerph14091010.
- Çelik A., Günay E., Aksu F., 2013. The evaluation of physical and motoric features of the elementary students between 7-9 age group. *Dokuz Eylül University Medical Faculty Journal*. 27(1), 7-13.
- Dumith S.C., Ramires V.V., Souza M.A., Moraes D.S., Petry F.G., Oliveira E.S., Ramires S.V., Hallal P.C., 2010. Overweight/obesity and physical fitness among children and adolescents. *Journal of Physical Activity and Health* 7, 641-648.
- Duncan M.J., Braynt E., Stodden D., 2017. Low fundamental movement skill proficiency is associated with high bmi and body fatness in girls but not boys aged 6–11 years old. *Journal of Sports Sciences* 21, 2135–2141. <http://dx.doi.org/10.1080/02640414.2016.1258483>.
- Evrin R.B., Fryar C.D., Wang C.Y., Miller I.V., Ogden C.L., 2014. Strength and body weight in us children and adolescents. *Pediatrics* 3, 1-19. doi:10.1542/peds.2014-0794.

- Fjortoft I., Pedersen A.V., Sigmundsson H., Vereijken B., 2011. Measuring physical fitness in children who are 5 to 12 years old with a test battery that is functional and easy to administer. *American Physical Therapy Association* 91, 1087–1095.
- Haslofça E., Haslofça F., Kutlay E., 2011. The Relations Between Physical Suitability Parameters on The Children Between 9-10 Ages. *Sports Medicine Journal* 46, 67-76.
- Henrique R.S., Bustamante A.V., Freitas D.L., Tani G., Katzmarzyk P.T., Maia J.A., 2017. Tracking of gross motor coordination in Portuguese children. *Journal of Sports Sciences* 1-9. <http://dx.doi.org/10.1080/02640414.2017.1297534>.
- Henrique R.S., Re A.H.N., Stodden D.F., Franssen J.F., Campos C.M.C., Queiroz D.R., Cattuzzo M.T., 2016. Association between sports participation, motor competence and weight status: a longitudinal study. *Journal of Science and Medicine in Sport* 19, 825–829. <http://dx.doi.org/10.1016/j.jsams.2015.12.512>.
- Karim O.A., Ammar, A., Chtourou H., Wagner M., Schlenker L., Parish A., Gaber T., Hökelmann A., Bös K., 2015. A comparative study of physical fitness among Egyptian and German children aged between 6 and 10 years. *Advances in Physical Education* 5, 7-17. <http://dx.doi.org/10.4236/ape.2015.51002>.
- Kim C.I., Lee K.Y., 2016. The relationship between fundamental movement skills and body mass index in Korean preschool children. *European Early Childhood Education Research Journal* 6, 928–935. <http://dx.doi.org/10.1080/1350293X.2016.1239326>.
- Lad U.P., Satyanarayana P., Lad S.S., Sırı C.H.C., Kumari, N.R., 2013. A study on the correlation between the body mass index (BMI), the body fat percentage, the handgrip strength and the handgrip endurance in underweight, normal weight and overweight adolescents. *Journal of Clinical and Diagnostic Research* 7(1), 51-54. DOI: 10.7860/JCDR/2012/5026.2668.
- Liao H.K., 2016. Hand grip strength in low, medium, and high body mass index males and females. *Middle East J Rehabil Health* 1, 1-7. doi: 10.17795/mejrh-33860.
- Marmeleira J., Veiga G., Cansado H., Raimundo A., 2017. Relationship between motor proficiency and body composition in 6- to 10-year-old children. *Journal of Paediatrics and Child Health* 7, 1-6. doi:10.1111/jpc.13446.
- Mayorga D.V., Merino R.M., Garcia J.C.R., 2013. Validity of sit-and-reach with plantar flexion test in children aged 10-12 years. *International Journal of Medicine and Science of Physical Activity and Sport* 15(59), 577-591.
- Montalcini T., Ferro Y., Salvati M.A., Romeo S., Miniero R., Pujia A., 2016. Gender difference in handgrip strength of Italian children aged 9 to 10 years. *Italian Journal of Pediatrics* 16, 1-6. doi 10.1186/s13052-016-0226-y.
- Omar M.T., Alghadir A., Baker S.A., 2014. Norms for hand grip strength in children aged 6–12 years in Saudi Arabia. *Developmental Neuro Rehabilitation* 1-6. doi:10.3109/17518423.2014.967878.



- Özkoçak, V., 2018. Body composition of the children between 5-14 age group. *Turkish Studies Social Sciences* 13(10), 875-885. <http://dx.doi.org/10.7827>.
- Ranson R., Stratton G., Taylor S.R., 2015. Digit ratio (2D:4D) and physical fitness (eurofit test battery) in school children. *Early Human Development* 91, 327–331. <http://dx.doi.org/10.1016/j.earlhumdev.2015.03.005>.
- Román P.A., Castillo R.M., Zurita M.L., Sánchez J.S., Pinillos F.G., López D.M., 2016. Physical fitness in preschool children: association with sex, age and weight status. *Child Care Health and Development* 1-7. DOI:10.3305/nh.2015.32.4.9486.
- Sánchez C.C., Artero E.G., Concha F., Leyton B., Kain J., 2015. Anthropometric characteristics and physical fitness level in relation to body weight status in Chilean preschool children. *Nutr Hosp* 1, 346-353. doi:10.3305/nh.2015.32.1.9092.
- Santos J.R., Ruiz J.R., Cohen D.D., Montesinos J.L., Pínero J.C., 2015. Reliability and validity of tests to assess lower-body muscular power in children. *Journal of Strength and Conditioning Research* 8, 2277–2285.
- Santos M.A.M.D., Almeida M.B.D., Castro R.M.D., Katzmarzyk P.T., Maia J.A.R., Leandro C.G., 2014. Birthweight, body composition, and motor performance in 7- to 10-year-old children. *Developmental Medicine & Child Neurology* 470-475. doi: 10.1111/dmcn.12664.
- Siahkoughian M., Mahmoodi H., Salehi M., 2011. Relationship between fundamental movement skills and body mass index in 7-to-8 year-old children. *World Applied Sciences Journal* 9, 1354-1360.
- Souza M.A., Baptista C.R., Benedicto M.M., Pizzato T.M., Sverzut A.C., 2014. Normative data for hand grip strength in healthy children measured with abulb dynamometer: a cross-sectional study. *Society of Physiotherapy* 1-6. <http://dx.doi.org/10.1016/j.physio.2013.11.004>.
- Souza M.C.D., Chaves R.N.D., Lopes V.P., Malina R.M., Garganta R., Seabra A., Maia J., 2014. Motor coordination, activity, and fitness at 6 years of age relative to activity and fitness at 10 years of age. *Journal of Physical Activity and Health* 11, 1239-1247. <http://dx.doi.org/10.1123/jpah.2012-0137>.
- Urlu, Y., 2014. Analyzing physical activity levels of the children between 10-12 age group (Antalya example). PhD Thesis, University of Balıkesir.
- Zaqout M., Vyncke K., Moreno L.A., Etayo P.D., Lauria F., Molnar D., Lissner L., Hunsberger M., Veidebaum T., Tornaritis M., Reisch L.A., Bammann K.A., Sprengeler O., Ahrens W., Michels N., 2016. Determinant factors of physical fitness in European children. *Int J Public Health* 1-10. doi 10.1007/s00038-016-0811-2.

Creative Commons licensing terms

Authors will retain the copyright of their published articles agreeing that a Creative Commons Attribution 4.0 International License (CC BY 4.0) terms will be applied to their work. Under the terms of this license, no permission is required from the author(s) or publisher for members of the community to copy, distribute, transmit or adapt the article content, providing a proper, prominent and unambiguous attribution to the authors in a manner that makes clear that the materials are being reused under permission of a Creative Commons License. Views, opinions and conclusions expressed in this research article are views, opinions and conclusions of the author(s). Open Access Publishing Group and European Journal of Physical Education and Sport Science shall not be responsible or answerable for any loss, damage or liability caused in relation to/arising out of conflict of interests, copyright violations and inappropriate or inaccurate use of any kind content related or integrated on the research work. All the published works are meeting the Open Access Publishing requirements and can be freely accessed, shared, modified, distributed and used in educational, commercial and non-commercial purposes under a [Creative Commons attribution 4.0 International License \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/).