



DOES OBSESSION OF IRRATIONAL STACKING ANABOLIC STEROIDS WITH TRENBOLONE ACETATE OVER DECADES LEADS TO GENERAL, CARDIOVASCULAR OR SOCIAL DEVIATION PROBLEM IN YOUNG ADULTS, OR JUST BIGGEST MUSCLE MASS IS EQUAL HIGHEST SOCIAL REPUTATION IN GYM AND IS THIS ALL PRICE HEALTH WORTH? - CASE REPORT

Sid Solaković¹ⁱ

Mensur Vrcić²,

Ratko Pavlović³

¹Cardiovascular Surgery Clinic University of Sarajevo,
Clinic Vascular and Endovascular Surgery Department,
Bosnia and Herzegovina

²Faculty of Sport and Physical Education, University of Sarajevo,
Bosnia and Herzegovina

³Faculty Physical Education and Sport, University of East Sarajevo,
Bosnia and Herzegovina

Abstract:

Inconspicuous epidemic of anabolic steroids application among young recreational athletes contributes to a greater potential damaging of the overall cardiovascular and mental state of obsessed individuals. All of them turn to anabolic steroids without a rational concept and take them in big amounts with unbelievable irrational combinations (stacking) of anabolic steroids of wide spectrum and questionable quality from black market with the aim to satisfy their pathological obsession of subjectivity of muscle mass, strength and appearance. An examinee, 36 years old (with the height of 182 cm and the weight of 168 kg), recreational amateur power lifter comes to doctor because of the subjective hardships in the form of tiredness and occasional pain in chest area with exact location (on the level of sternum and pectoral muscles), denies retrosternal pain. Also, he reports dizziness and occasional leg pain. He suspects that he has insufficiency of artery peripheral circulation. Personally, he also names frequent insomnia, decrease of tolerance levels alongside tolerated aggressiveness and anger attacks with auto mutilation and periodical depression

Key words: anabolic steroids, cardiovascular problem, social deviation

ⁱ Correspondence: email sid.solakovic@gmail.com

1. Introduction

The biggest muscles at all costs and the question whether they are worth the health are a game of Russian roulette in each individual who uncontrollably and irrationally uses excessive amounts of anabolic steroids. Quiet epidemic on an alarming level of potential cardiovascular dangers as well as social deviant behaviour manifested in the form of uncontrollable aggression, depression and physical violence among young population of recreational bodybuilding, fitness and power lifting population has become a subject of serious medical-social debate. ^{1-5,11} Limitation of research on consequences of isolated anabolic medications of today's world is in constant correlation because of the subjective effects of stacking doses among many individuals. Also, restraint is a consequence because of the saturation from various medications on the black market. There are differences in terms of medication quality as well as many legal and illegal pharmacological manufacturers and inability to oversee them in a long term sense. There is certain secrecy in terms of consuming during doctor's appointments, as well as the frequency of appointments themselves, which limits the access to adequate medical science data, the damage to endothelial function as well as cerebral cardiovascular system and progress level of damage to peripheral circulation. ⁵⁻¹⁴ The question is whether the problem of epidemic obsessive consummation among young recreational users is serious and is multi discipline medical-socio-psychological therapy treatment is needed, or is it just the case for medical treatment with permanent social deviation for unreal physical super performances?

2. Case Report

36 year-old recreational power lifter with occasional periods of amateur performances on competitions comes to an emergency medical centre because of subjective hardships of tiredness and occasional chest pain (on the level of pectoral muscles), denies retrosternal pain with propagation to neck, head and upper extremities. Also, he names episodes of occasional pressure in the head, vertigo and emotional irritation and doubts to a brain stroke. In the information from the anamnesis he also names frequent insomnia episodes, lowered tolerance threshold with heightened aggressiveness and anger episodes with periodical deep depression. He names the start of oral and intramuscular application of anabolic steroids from the age of 18 (13 years of continuous consuming with occasional pauses of 2 month cleanses depending on the stacking). Also, he says that after consummation of Trenbolone acetate (weekly dose), after three months of application he has frequent need to self-injure alongside enormous subjectivity and objectivity of increase of muscle mass and increase of maximum and sub maximum strength, which disappear after they stop using anabolic substances (Trenbolone acetate) in comparison to other steroids in stacking. In anamnestic data he denies lack of control and aggressive episodes as a positive family

DOES OBSESSION OF IRRATIONAL STACKING ANABOLIC STEROIDS WITH TRENBOLONE ACETATE
OVER DECADES LEADS TO GENERAL, CARDIOVASCULAR OR SOCIAL DEVIATION PROBLEM IN YOUNG
ADULTS, OR JUST BIGGEST MUSCLE MASS IS EQUAL HIGHEST SOCIAL REPUTATION
IN GYM AND IS THIS ALL PRICE HEALTH WORTH? - CASE REPORT

anamnesis of mental illnesses. The question here is whether Trenbolone acetate and Trenbolone Hexahydrobenzylcarbonate is so aggressive and destructive steroid type and does it have any connection to aggressiveness and deviant social behaviour, bearing in mind the already applied irrational stacking which the patient has used for years.



Figure 1*

Figure 2*

Figure 3*

Figure 2 and Figure 3, Automutilization on the left upper arm after 3 months Trenbolone acetate in agreement with other anabolic preparations

Table 1: Parameters and values of the power lifter men

Age	36
High	182cm
Weight	168kg
Systolic blood pressure	165 (mmHg)
Diastolic blood pressure	90 (mmHg)
Left ACC Carotid Intima-Media Thickness CIMT(mm)	0,90 ($\pm 0,02$)
Right ACC Carotid Intima-Media Thickness CIMT(mm)	0,93($\pm 0,04$)
Left Internal Carotid artery PSV 95 cm/sec	-
Left Internal Carotid artery EDV35 cm/sec Left ICA/CCA PSV ratio <2.0	-
Right InternalCarotid artery PSV85 cm/sec	-
Right Internal Carotid artery PSV32 cm/sec	-
Right ICA/CCA PSV ratio <2.0	-
Presense of atheromatose plaques on carotid bifurcation arteries	yes
Triglycerids	2,4
LDL low-density lipoprotein Cholesterol	4,50 mmol/L
HDL high-density lipoprotein Cholesterol	0,79 mmol/L
Total Cholesterol	7,0 mmol/L
RBC (eritrociti)	5,32 x 10 ¹² / L
HGB (hemoglobin)	182 g/L
HCT (haematokrit)	0.57 l/L
PLT (trombociti)	169/L

*from author's collection.

2.1 Oral and intramuscular application of continuously anabolic steroids approximately over 12 years with cleaning period 2-6 months in each year

Table 2: An individual stacking concept was applied without the concept of the sequence of anabolic substance applications

Currently consumed (Methandrostenolone)	25-75mg oral.aplication
Currently consumed (Oxandrolone)	30mg-50mg oral application
Currently consumed (Stanozolol)	intramuscular aplication
Currently consumed (Stanozolol)	10 miligram / oral aplication 3 time
Currently consumed Nandrolone Decanoate	400ml per week i.m
Currently consumed Testosterone propionate	1ml every 2 days i.m
Currently consumed Trenbolone acetate	75-100mg oral
Currently consumed testo mix 4 form of testesteron	1,5 ml per week i.m
Currently consumed Testosterone Enatate (depo)	1ml per week i.m
Currently consumed Drostanolone propionate	1ml per week i.m
Currently consumed Trenbolone Hexahydrobenzylcarbonate	1ml per week i.m



Figure 4: Current anabolic supplementation used by the patientⁱⁱ

3. Discussion

Automutilation is defined as making injuries to oneself without suicidal intents. Although many studies of anabolic steroids mostly are focused on the appearance of aggressiveness towards other people, as well as criminal, asocial, delinquent behaviour, the scientific facts about auto mutilation during the consumption of anabolic medications are lesser known, so the question is whether in very big doses, even during excessive stacking, can start such a type of individual asocial behaviour?^{1-5,11} Various studies defined the misuse of anabolic steroids through aggressive episodes and behaviour disorders. Also, very often, potential users exchange the use of anabolic steroids with other drug, such as heroin or cannabis consuming.^{6,7} On the other hand,

ⁱⁱ from author's collection.

the science proved that stacking (using many different drugs) increases muscle performance and thus the overall performance is improved. Some studies were conducted regarding the potential damage of synthetic testosterone and other types of derivatives of synthetic testosterone, which after all leads to deviant behaviour. Application doses in misuse of anabolic steroids among weight lifters and bodybuilders outperform medical dosages in 10 and up to 50 times of recommended medical dose for the patients. Using of anabolic steroids usually starts in cycles of 2 steroids which lowers to previously irrational limit of subjectivity of combination effect of the so called "stacking" (increase and influence of several drugs) because of the effect in cycles of 3-6 months depending on the potential of anabolic substance itself with cleansing pauses.⁷⁻¹⁵

Accommodation cycles start with a small application dose, usually orally or intramuscular, which increases because of the subjective expectations of training quality, and decreases after possible cycle termination. Bearing in mind that in the periods of cleanse a sudden decrease occurs in terms of subjective strength estimation, some individuals who are inexperienced, exchange drugs with other substances because they think it saturates the receptors and effectiveness of the forbidden substance. Anabolic steroids are specific and they are associated with a high number of cerebrovascular and heart incidents worldwide among younger population under the age of 30, while frequent incidents are recorded among the population of the age over 40, and usually among individuals who are long-term users.¹⁵⁻¹⁸ Because of the limitation of the number of examinees, as well as the studies themselves, and secrecy about the consumption among research subjects, it is impossible to determine which stacking combination is associated with the highest possibility of atherosclerosis illness, cerebrovascular diseases, in terms of obstruction of artery and vein hemodynamic because of the changes on the level of total cholesterol, triglycerides as well as direct impact on the decrease of HDL and increase of the LDL.

In the 50s and late 70s anabolic steroids in certain dosages do not make the process faster, and have therapeutic effect in terms of arteriosclerosis process prevention with myocardial infarction and symptomatology of chronic stenosis occlusive artery illness. It is determined that except individual subjective behaviour, there is a minimal effect of increase and disorder of lipid status in terms of decrease of HDL and LDL increase. Studies which did research on long-term users of anabolic steroids, which surpassed all rational doses, negatively report on that.¹⁸⁻²²

Having in mind that the subject of the research is sports and now widespread consumption of epidemic levels in recreational purposes, drugs manipulation, and the role is not clear in terms of preventive therapy effects on cardiovascular system and potential damage of various doses and various substance combinations. A field that has not been yet researched thoroughly is among the population of the age between 20 and 30. The science still does not have available information and is limited in regards to the numbers of such research. Further research is needed of secondary and primary

prevention of arteriomatosis process, as well as general suppression of arteriosclerosis in risky amount of young recreational body building population.²²⁻³¹

Studies of medicine applications of Stanazolol and Ester Nandrolon are scientifically significantly resulted in inadequate application on delevation of lipid level in serum and long-term misuse of supra-therapeutic doses of such substances carry a highly dangerous potential of cardiovascular incident. Therapeutically overdosed misuse of anabolic steroids has impact on decrease of HDL and LDL increase and provokes disorder of lipids of high and low density relation (HDL and LDL), which leads to the start of arteriomatosis with the progression of adhesion of thrombocyte and speed increase of the pathology of atherosclerosis because of increased values of erythrocytes and thrombocytes as well as haematocrits. The sole backing of the research works and case report is a statistical indicator of irrational epidemic increase tendency of sole application of anabolic steroid in terms of drug usage in young recreational body building Balkan population. It is evident that without the usage of anabolic some recreational athletes do not even think about starting a body building training process and by using irrational stacking applications cardiovascular incidents after long-term usage could become more frequent.^{12,22-29}

Contemporary recreational group of anabolic substances users is greater in numbers than the starting group from 1970 and 1980 and is far more visible and presented with the public. We can even talk about the starting euphoria of obsession and Addison complex with the development of epidemic scales. The scale is such that application doses of synthetic testosterone are going over irrational limits of thinking, where among some individuals they range between 3000-5000 mg several times in a week alongside application doses of various anabolic steroids simultaneously, which has no benefit from application besides toxic properties and occurrence of deviant personality due to hormonal misbalance of the individual. Many studies led to conclusion that the problem of the start of consuming has different roots (enormous strength with social respect, imagined self-confidence gain, imagined superiority) and that there is a potential of self-harm, harming other people or a murder. From our standpoint more attention should be given to the population which is included in social, cardiovascular, psychiatric as well as potential criminal risky population. Several questions can be asked. Is there only irrational consumption of anabolic substances or does the genetic predisposition with gained form of social deviant behaviour has any impact? Is one of the anabolic substances, eg. Trenbolon acetate or hexahydrobenzylcarbonate the key to the initial change of personality or is it created by some other substance combinations? Many other studies will be needed in order for this serious topic of the dark side of sport to be researched accordingly.^{2,4,5,27,28,29}

Overall a lot of efforts and many researches will be necessary to gain precise information on usage and application of anabolic steroids in order to come to an exact initiator of irrational application and in order for medicine to act not only

symptomatically and therapeutically but preventively and cardioprotectively as well.^{2,4,25}

4. Conclusion and Further Research

Epidemic increase of usage of anabolic steroids has been recorded among young population and the number keeps increasing. The obsession for irrational use of anabolic steroids is a serious problem of addiction equivalent to using drugs with bad consequences on cardiovascular system, deviant social behaviour as well as strong deviant factor of social domination among male body building recreational population. Irrational overdosed and continuous stacking of anabolic substances with the subjective aims to increase strength and mass leaves a strong psychological and social deviant form of behaviour which has a strong destructive cardiovascular potential with the personification disorder as a strong psychiatric, and in some cases self-harming potential. Overall, we can speak about a new young cardiovascular risky population which has not shown itself to the public yet, and represents a big problem for a wider social environment

Conflict of Interest

The authors declare no conflict of interest

References

1. Petersson A., Benqtsson J., Voltaire-Carlsson A., & Thiblin I. (2010). Substance abusers' motives for using anabolic androgenic steroids. *Drug Alcohol Depend* (111), 170–172.
2. Beaver K.M., Vaughn M.G., DeLisi M., & Wright J.P. (2008). Anabolic-androgenic steroid use and involvement in violent behavior in a nationally representative sample of young adult males in the United States. *Am J Public Health* (98), 2185–2186.
3. Midgley S.J., Heather N., & Davies J.B. (2001). Levels of aggression among a group of anabolic-androgenic steroid users. *Med Sci Law* (41), 309–314.
4. Lundholm L., Frisell T., Lichtenstein P., & Langstrom N. (2015). Anabolic androgenic steroids and violent offending: confounding by polysubstance abuse among 10 365 general population men. *Addiction* (110), 100–108.
5. Choi P.Y., & Pope H.G. Jr. (1994). Violence toward women and illicit androgenic-anabolic steroid use. *Ann Clin Psychiatry* (6), 21–25.
6. National Institute on Drug Abuse Research Report Series: Anabolic Steroid Abuse, 2006.

7. Schulze, Jenny J. & et.al. (2008). Doping Test Results Dependent on Genotype of UGT2B 17, the Major Enzyme for Testosterone Glucuronidation. *Journal of Clinical Endocrinology & Metabolism* March 11, 2018.
8. King D.S., Sharp R.L., & Vukovic M.D. (1999). Effect of oral androstenedione on serum testosterone and adaptations to resistance training in young men: a randomized controlled trial. *J Am Med Assn.* (281), 2020-2028.
9. Wilson J.D. (1988). Androgen abuse by athletes. *Endocr Rev.* (9), 181-191.
10. Yeater R., Reed C., Ullrich I. & et al. (1996). Resistance trained athletes using or not using anabolic steroids compared to runners: effects on cardio respiratory variables, body composition, plasma lipids. *Br J Sports Med.* (30), 11-14.
11. Muller R.W., & Hollmann W. (1998). Akute Lipoproteibeeinflussung durchein anaboles Steroid bei Kraftsportlern. *Dtsch Z Sportmed.* (39), 35-40.
12. Zmuda J.M., Fahrenbach M.C., Younkin B.T., & et al. (1993). The effect of testosterone aromatization on high-density lipoprotein cholesterol level and postheparin lipolytic activity. *Metabolism* (42), 446-450.
13. Ale'n M., &Suominen J. (1984). Effect of androgenic and anabolic steroids on spermatogenesis in power athletes. *Int J Sports Med.* (5), 189-192.
14. Durstine J.L., & Haskell W.L. (1994). Effects of exercise training on plasma lipids and lipoproteins. In: Holloszy JO, editor. Exercise and sport sciences reviews. Baltimore (MD): Williams & Wilkins, 477-522.
15. Applebaum-Bowden D., Haffner S.M., & Hazzard WR. (1987). The dyslipoproteinemia of anabolic steroid therapy: increase in hepatic triglyceride lipase precedes the decrease in high density lipoprotein cholesterol. *Metabolism* (36), 949-952.
16. Taggart H.M., Applebaum B.D., Haffner S., & et al. (1982). Reduction in high density lipoproteins by anabolic steroid (stanozolol) therapy for postmenopausal osteoporosis. *Metabolism* (31), 1147-1152.
17. Turillazzi, E., Perilli, G., Di Paolo, M., Neri, M., Riezzo, I. & Fineschi, V. (2011). Side effects of AAS abuse: an overview. *Mini Rev. Med. Chem.* (11), 374-389.
18. Severo, C., Ribeiro, J., Umpierre, D., Silveira, A.D., Padilha, M., De Aquino Neto, F. & Stein, R. (2012) Increased atherothrombotic markers and endothelial dysfunction in steroid users. *Eur. J. Prev. Cardiol.* 20 (2), 195e201.
19. Fernandez-Balsells, M. Murad, M. Lane, J.F. Lampropulos, F. Albuquerque, R.J. Mullan, N. Agrwal, M. Elamin, J. Gallegos-Orozco, A. Wang, P. Erwin, S. Bhasin, V. Montori (2010). Clinical review 1: adverse effects of testosterone therapy in adult men: a systematic review and meta-analysis. *J. Clin. Endocrinol. Metab.* 95 (6), 2560-2575.
20. Dawson, R. (2001). Drugs in sport-the role of the physician. *J. Endocrinol.* 170 (1), 55-61.
21. Ammatuna, E., Nijziel, M. (2014). Polycythemia and renal infarction in a bodybuilder. *Q. J. Med.* 107, 661.

22. Shamsavarinia, K., Rahmani, F., Ebrahimi Bakhtavar, F., Hashemi Aghdam, Y., Balafar, M. (2014). A young man with myocardial infection due to trenbolone acetate; a case report. *Emergency* 2 (1), 43-45.
23. Hartgens, F., Kuipers, H. & et al. (2004). Effects of androgenic-anabolic steroids in athletes. *Sports Med.* 34(8), 513-54.
24. Evans N.A. et al. (2006). Current concepts in anabolic-androgenic steroids. *Am J Sports Med.* 32(2), 534-542.
25. Kindermann W. (2006). Cardiovascular side effects of anabolic-androgenic steroids. *Herz* 31(6), 566-573.
26. Haug, E. et al., Androgenic-Anabolic Steroids (AAS) and Violent Behaviour. Report from Norwegian Knowledge Centre for the Health Services (NOKC) No. 04-200.
27. Laye-Gindhu, A., Schonert-Reichl, Kimberly A. (2005). "Nonsuicidal Self-Harm Among Community Adolescents: Understanding the "Whats" and "Whys" of Self-Harm". *Journal of Youth and Adolescence.* 34 (5), 447-457.
28. Klonsky, D. (2007). The functions of deliberate self-injury: A review of the evidence. *Clinical Psychological Review.* 27 (2), 226-239.
29. Muehlenkamp, J.J. (2005). Self-Injurious Behavior as a Separate Clinical Syndrome. *American Journal of Orthopsychiatry.* 75 (2), 324-333.

Sid Solaković, Mensur Vrcić, Ratko Pavlović
DOES OBSESSION OF IRRATIONAL STACKING ANABOLIC STEROIDS WITH TRENBOLONE ACETATE
OVER DECADES LEADS TO GENERAL, CARDIOVASCULAR OR SOCIAL DEVIATION PROBLEM IN YOUNG
ADULTS, OR JUST BIGGEST MUSCLE MASS IS EQUAL HIGHEST SOCIAL REPUTATION
IN GYM AND IS THIS ALL PRICE HEALTH WORTH? - CASE REPORT

Creative Commons licensing terms

Authors will retain the copyright of their published articles agreeing that a Creative Commons Attribution 4.0 International License (CC BY 4.0) terms will be applied to their work. Under the terms of this license, no permission is required from the author(s) or publisher for members of the community to copy, distribute, transmit or adapt the article content, providing a proper, prominent and unambiguous attribution to the authors in a manner that makes clear that the materials are being reused under permission of a Creative Commons License. Views, opinions and conclusions expressed in this research article are views, opinions and conclusions of the author(s). Open Access Publishing Group and European Journal of Physical Education and Sport Science shall not be responsible or answerable for any loss, damage or liability caused in relation to/arising out of conflict of interests, copyright violations and inappropriate or inaccurate use of any kind content related or integrated on the research work. All the published works are meeting the Open Access Publishing requirements and can be freely accessed, shared, modified, distributed and used in educational, commercial and non-commercial purposes under a [Creative Commons attribution 4.0 International License \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/).