



THE EFFECTS OF SIX-WEEK AEROBIC EXERCISE PROGRAM ON BODY COMPOSITION AND BLOOD LIPIDS IN WOMEN

Muzaffer Selçuk¹, H. Bayram Temur¹ⁱ,

Salih Öner², Eda Kinaci²

¹Yüzüncü Yıl University,
Physical Education and Sports High School,
Van, Turkey

²Firat University,
Sports Sciences Faculty,
Elazığ, Turkey

Abstract:

Introduction and Purpose: This study was conducted to investigate the effects of 6 Week Aerobic Exercise Program on Body Composition and Blood Lipids in Women.

Methods: The population of the research consists of women who make step aerobics in the fitness centers of Elazığ and the sample consists in total 12 sedentary women of between the ages of 25 and 30 selected by random method. Participants were given an aerobic exercise program for 3 days and 30 minutes a week. The intensity of exercise was determined by Karvonen method as 50-60%. Participants' body weights, body fat percentages and blood lipids were measured before and after exercise. In the analysis of the data, SPSS 17 package program was used. The obtained data were evaluated statistically at ($p < 0.05$) level.

Findings: There was a significant difference in body weight and body fat percentages of sedentary women participating in the study compared to before exercise ($p < 0.05$). When the blood lipids of the participants were examined, significant differences were found in HDL LDL and triglyceride levels compared to pre-test values ($p < 0.05$).

Discussion and Conclusion: As a result, regular aerobic and anaerobic positive effect on body composition and blood lipids in women aged 25-30. It can be said that such regular training is necessary for a healthy life, especially for the prevention of obesity.

ⁱ Correspondence: email temurb@hotmail.com

Keywords: women, exercise, blood lipids, physical fitness

Introduction

In addition to technological developments in our time, new living conditions lead people to a still, still life. When we look at the World Health Organization's 2002 review, still life, in other words, static life causes the world to lose 1.8 million lives every year. In the world, inactivity affects about 11-16% of breast, colon cancer and diabetes mellitus and 23% of heart disease (Akyol et al., 2008). Constantly, changing technology forces individuals to do less physical activity in their living space. With the further development of computer games in the 1970s, recreational activities for people of all ages have entered a new era (Turkish Public Health Institution).

Today's fast-paced technology has reduced the need for human beings, and the response from the work environment, which is a life model opposite to the natural structure of the human being, has entered into death causes, especially the disturbances of the stress digestive system. Sports is a solution against this danger placed against the modern people by developing a field that is far away from the troubles brought by daily life and also helping medicine by the healthy life style that it brings. This benefit has positive effects on the social development of the people. Therefore, the developed countries place more emphasis on spore and give physical education and sports lessons to children starting from early ages (Gökhan et al., 1979). Increasing people's awareness, level of exercise and health make people's life better. In childhood and youth, making physical activity a part of life will ensure a healthy and more peaceful life in the next life. In this sense, knowing what to do for women to have a healthy lifestyle will benefit people in terms of better quality of life (Ökan A., 2013).

Exercisers are expected to have a number of physiological changes with both acute and chronic adaptation. Regular long-term and moderate-intensity aerobic exercises are reported to increase the level of high-density lipoprotein (HDL-C), which reduces lipids such as Total Cholesterol, LDL-C and Triglyceride, which are risk factors for coronary artery disease. It is also emphasized that high blood pressure and obesity diseases decrease with exercise (Akgün N., 1995, Fox 1999, Çolakoğlu 2003.). However, the effect of doing exercises on some blood samples just as thyroid hormones depends on gender, datum fitness level, nutritional status, environmental temperature, height, the intensity and type of the exercise and the time measurement is done (Bernet and Wartofsky 2000, Polat et al 2017).

2. Materials and Methods

This study is composed of women performing step aerobics in the fitness centers of Elazığ and their sample is composed of 12 sedentary women between 25 and 30 years old selected by random method. Participants were asked to have a doctor's report that they did not mind any exercise. Blood lipids (Total-C, HDL-C, LDL-C, and triglyceride) were measured by age, body weight, resting heart rate, body fat percentage. For this, an average of 7 cc fasting blood samples was taken.

2.1 Exercise Program

Athletes were subjected to a run-and-walk exercise at 50-60% of the target heart rate, 6 weeks, 3 days a week, 30 minutes according to the progressive training method. Subjects were given a 10 min warm-up exercise before starting the workout, and a 10 min stretching exercise at the end of the workout.

The intensity of the exercise was determined immediately after the end of exercise by the number of target heart rate according to the number of heartbeats (Karvonen) method as a result of the 10 second heart rate count from the carotid artery in the neck (Fox 1990, Tamer 2000, Zorba 1995.).

The method is shown below:

$$\text{HRmax} = 220 - \text{years}, \text{HRR} = \text{HR max} - \text{HR rest}, 60\% \text{ THR} = (0.60 \times \text{HRR}) + \text{HR rest}.$$

All measurements and tests applied to the participant were performed twice, one week before the training program (pre-test) and one week after the training program (post-test).

2.3 Statistical Analysis

SPSS 17 package program was used in the analysis of the data. The obtained data were analyzed by standard deviation, averaged paired comparison analysis, Paired Samples T Test, statistically ($p < 0.05$).

3. Results

Table 1: Participants' Demographic Information

Variables	Person Num	%	mean	Standard deviation	Minimum	Maximum
Age	12	100	27,5	1,78	25	30
Height	12	100	1,65	0,04	1,58	1,73

Table 2: Analysis of Participants' Pre-Test and Post-Test Values

Variables		Person Num	mean	Standard deviation	P
Weight	Pre-Test	12	57,5	4,03	0,00
	Post-Test	12	54,4	3,70	
Body Fat Percentage	Pre-Test	12	26,7	5,50	0,02
	Post-Test	12	21,2	3,30	
Body Mass Index	Pre-Test	12	21,1	1,12	0,00
	Post-Test	12	19,9	1,02	
Hdl	Pre-Test	12	41,8	2,79	0,01
	Post-Test	12	45,0	2,90	
Ldl	Pre-Test	12	83,8	14,6	0,00
	Post-Test	12	72,0	11,4	
Triglycerides	Pre-Test	12	199,5	81,7	0,01
	Post-Test	12	131,7	48,3	

Table 2 shows that ; there is statistically significant differences in participants' weight, body fat percentage, body mass index, HDL, LDL, and triglyceride pre-test and post-test values.(p <0,05).

4. Discussion and Conclusion

Pre-test averages of 57,5 kg were determined for women who have step aerobics between 25-30 years of age participating in the study, while post-test averages were 54,4 kg. Body fat percentage in pre-test and post-test measurements was found to be decreased in step aerobics for body mass index measurements. Increase in HDL values in blood lipids decreased in LDL values. It is thought that 6 weeks training is effective on Body Composition and Blood Lipids.

Babayiğit et al. (2002), found that sedentary women aged 25-31 years had a decrease in body weight at the end of the 8-week, 3-week, medium-severity 45-minute step program.

Çolakoğlu and Karaca (2006) found that body weight decreased in both groups when they applied aerobic exercise for 12- week, 3-days 45-60 minutes and 50-75%

intensity to middle-aged and young women. In another study, Çınar et al. (2017) found that six-week weight lifting exercises reduced body weight.

In a study by Fatouros et al. (2002) 32-person group of ages between 65 and 78 years reported a reduction in body weights in a 16-week, 3-week, 55-80% aerobic exercise trial.

In the study performed by Imamoğlu et al. (2002), on 45 sedentary it has been found that there was no significant decrease in body weight, physical fitness, body composition and some blood parameters of three-day, one-hour, three-month exercise three times a week in the study that they made.

Yaprak (2004), found a significant decrease in body weights and body mass indexes in the study he made on 41 obese women which, aged $37,70 \pm 2,75$ years, $158,64 \pm 5,59$ cm in length and $83,12 \pm 11,56$ kg, who do 45-60 minutes of aerobic exercise per day and three days in a week eight weeks.

In the study performed by Çolakoğlu and Karacan, (2006) 8-week aerobic exercise was found to decrease in body mass index at the end of 3-day 30-minute run-walk exercise program for 12 weeks.

Body fat percentage of the study group was measured as $27,566 \pm 5,070\%$ before exercise program and $25,613 \pm 5,054\%$ after exercise program. The study group's pre-exercise program values and post-exercise program values were found to be significantly decreased after the exercise program (Kurt 2007).

Murphy and Hardman (1998) reported that, there was a significant decrease in body fat percentages of 47 sedentary that performed short and long fast walks in the study that they organized as; first group walk 10 minutes each day and second group 30 min.

As a result of the study that made by Ug et al. (2006) it has been reported that the body fat ratio decreases in the subject group who do regular endurance type of exercises not less than 3 hours a week, between 78 men and 53 women aged 20-74 years. Özdal et al. (2016) reported that the practice of Body Vibration exercise reduced the fat percentage of the research group.

As a result of other studies on body fat percentage, it has been reported that body fat percentage decreases (Murphy and Hardman 1998, Sevim et al. 1996, Tsourlou et al. 2003).

Dönmez and Aydos (2000) examined the changes in blood lipids and body fat mass with 20-week aerobic exercise in a total of 650 male and female subjects aged between 17 and 65 years. At the end of the training, subjects had a 3.3% decrease in body fat mass and a significant correlation was found between changes in body fat mass, and LDL-C, total cholesterol, total-K / HDL-C lipid exchange index of women.

A number of studies have been conducted on the effects of aerobic exercises on physical fitness and blood lipids (Dönmez and Aydos 2000, Karacan and Çolakoğlu 2003, Çınar et al. 2016).

In particular, the number of studies reporting that regular and controlled exercises reduce body fat and increase lean body mass are increasing day by day (Çınar et al. 2016, Pancar e et al. 2017).

As a result, six weeks of step aerobic exercises have a positive effect on weight, body fat percentage, body mass index, HDL, LDL, Triglyceride variables. Six-week step-aerobic exercise can be applied to sedentary women aged 25-30 years. Different durations and different exercise models can compare the effects of the same age and different age groups.

References

1. Akgün, N.; "Place of Exercise in the Treatment of Obesity", *Journal of Sports Medicine*, 20(3):91-103, 1995.
2. Akyol A, Bilgiç P, Ersoy, G. *Physical Activity, Nutrition and Healthy Living*. Hacettepe University-Faculty of Health Sciences Department of Nutrition and Dietetics, Ankara, 2008.
3. Babayiğit, G., Zorba, E., İrez, S.G. Ve Mollaoğullari, H.(2002). "The Effect of 8 Week Step Studies on Some Physiological and Anthropometric Values in 25-31 Years Old Women." 7th International Sports Science Congress. 27-29 Oct. 2002, s.156.
4. Bernet VJ, Wartofsky L. Thyroid function and exercise. *Contemp Endocrinol Sports Endocrinol* 2000.
5. Cinar V, Akbulut T, Sarikaya M. Effect of Zinc Supplement and Weight Lifting Exercise on Thyroid Hormone Levels. *Indian J Physiol Pharmacol* 2017; 61(3) : 232–236
6. Çınar V. Akbulut T. Öner S. Pancar Z. Karaman ME. "An Investigation Of Healty Life Style Behaviors of Turkish Wrestling Federation Coaches", *International Refereed Academic Journal of Sports, Health and Medical Sciences*, 21: 119-136, 2016.
7. Çolakoğlu F at al., "Effects of Eight-Week Aerobic Exercise Program on Sedentary Middle-Aged Women's Body Composition and Blood Lipids" *Gazi University School of Physical Education and Sports*, 2003.

8. Çolakoğlu, F.F. "The Effect of 8-Week Run-Walk Exercise on Physiology, Motoric and Somatotype Values of Sedentary Middle-Aged Obese Women." Gazi University, Journal of Gazi Education Faculty, 23(3):275-290. (2003).
9. Çolakoğlu, F.F. ve Karacan, S. "The Effect of Aerobic Exercise on Some Physiological Parameters in Young Women and Middle - Aged Women." Kastamonu Education Faculty, 14(1):277-284. (2006).
10. Dönmez G, Aydos L, The Effect of Callisthenic Workouts on the Physiological and Physical Parameters of Middle-aged Sedentary Women, Journal of Gazi Physical Education and Sport Sciences, 2: 17 – 25, Ankara, 2000
11. Fatouros, Ig., Taxildaris, K., Tokmakidis Sp., Kalapotharakos, V., Aggelousis, N., Athanasopoulos, S., Zeeris I. Ve Katrabasas I. (2002). "The effect of Strength Training, Cardiovascular Training and Their Combination on Flexibility of Inactive Olders Adults." Int J Sports Med, 23(2):112-119.
12. Fox, Bowers, Foss; "Physiological Foundations of Physical Education and Sports", Translation: Cerit, M., Bağırhan Publishing, Ankara. 1999.
13. Gökhan N, Olgun P, Gürses Ç. Sportive Ability Research Method Turkey Apptlication, TSV, İstanbul,2,1979
14. İmamoğlu, O., Akyol, P. Ve Bayram, L. (2002). "The Effect of Three Months Exercise on Physical Fitness, Body Composition and Some Blood Parameters in Sedanter Women." 7th International Sports Science Congress. 27-29 Oct 2002, s.19.
15. Karacan S, Çolakoğlu F, The Effect of Aerobic Exercise on Body Composition and Blood Lipids in Middle Aged Women and Young Sedentary Women., Spormetre Journal of Physical Education and Sports Sciences, 1(2) 83-88, 2003.
16. Kurt S. To investigate the effect of 8-week step-aerobic exercise program on some physical fitness parameters. 2007.
17. Murphy, MH. ve Hardman, AE. "Training Effects of Short and Long Bouts of Brisk Wolking in Sedentary Women." Med Sci Sports Exerc., 30(1):152-157. (1998).
18. Ökan A., Bozkuş T., Kul M., Türkmen M., Öz Ü., Cengiz C., "The Determination and Correlations of Physical Activity Levels and Healthy Life Style Behaviors of Public Performers. "International Journal of Science Culture and Sport (Int J SCS). September 2013; 1-3
19. Özdal M, Çınar V, Bostancı Ö, Kabadayı M, Akcan F. Effect of whole-body vibration training on body fat mass of sedentary healthy women .International Journal of Sport Studies, 6(1)2016

20. Pancar Z. Özdal M. Çınar V., "The Effect Of 4-Weekly Low Intensity Physical Activity Program In Thyroid Hormone Levels In Obese And Overweight Children", *European Journal of Physical Education and Sport Science*, 11(3):1-8,2017.
21. Polat M, Polat Y, Akbulut T, Cinar V, Marangoz I, The effects of trainings applied with CoQ10 and zinc supplementation on the thyroid hormone metabolism in Soccer players. *Biomedical Research* 2017; 28 (16): 7070-7075.
22. Sevim, M., Sevim, Y., Günay, M. VE Erol, E. "An Analysis of the Effect of Combined Strength Training on Performance Development of 18-25 Years Group Elite Female Handball Players." *Journal of Gazi Physical Education and Sport Sciences*, 1(3):2-7. (1996).
23. Tamer, K.; "Measurement and Evaluation of Physical - Physiological Performance in Sports", Bağırhan Publishing, Ankara-2000.
24. Tsourlou, T., Gerodimos, V., Kellis, E., Stavropoulos, N. ve Kellis, S. "The Effects of a Calisthenics and a Light Strength Training Program on Lower Limb Muscle Strength and Body Composition in Mature Women." *J Strength Cond Res.*, 17(3):590-598. (2003).
25. Turkish Public Health Institution. Access: (<http://fizikselaktivite.gov.tr/tr/yetiskinlerde-fiziksel-aktivite/isyerinde-fiziksel-aktivite-onerileri/>) Access date: 08.04.2015.
26. Yaprak, Y. "Effects of Aerobic and Strength Workouts on Oxygen Utilization and Heart Debate in Obese Women." *Spormetre. Journal of Physical Education and Sports Sciences*, 2(2):73-80. (2004).
27. Zorba, E., Ziyagil, M.A.; "Body Composition and Measurement Methods", Baskı; Erek Ofset, Trabzon -1995.

Creative Commons licensing terms

Author(s) will retain the copyright of their published articles agreeing that a Creative Commons Attribution 4.0 International License (CC BY 4.0) terms will be applied to their work. Under the terms of this license, no permission is required from the author(s) or publisher for members of the community to copy, distribute, transmit or adapt the article content, providing a proper, prominent and unambiguous attribution to the authors in a manner that makes clear that the materials are being reused under permission of a Creative Commons License. Views, opinions and conclusions expressed in this research article are views, opinions and conclusions of the author(s). Open Access Publishing Group and European Journal of Education Studies shall not be responsible or answerable for any loss, damage or liability caused in relation to/arising out of conflicts of interest, copyright violations and inappropriate or inaccurate use of any kind content related or integrated into the research work. All the published works are meeting the Open Access Publishing requirements and can be freely accessed, shared, modified, distributed and used in educational, commercial and non-commercial purposes under a [Creative Commons Attribution 4.0 International License \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/).